

A Rare Case of Masquerading Chondrosarcoma with Pathological Fracture of Neck of Femur – A Case Report

Ananya Mishra¹, Pulkit Gupta², Manohar³, Hoysala Kantharaju³, Dhyaan Bhardwaj³, Vipul Jhajharia³

Learning Point of the Article:

Be open for surprises in cases of tumors. Always send pre-operative and post-operative biopsy to confirm your diagnosis and for further treatment.

Abstract

Introduction: Chondrosarcoma is the second most common primary malignant tumor accounting for 20–27% of these. These are malignant cartilaginous tumor found most commonly in the pelvis, followed by the proximal femur. Perivascular epithelioid cell tumor (PEComa) is a rare mesenchymal neoplasm. These tumors all share a distinctive cell type, the perivascular epithelioid cell (which has no known normal tissue counter-part).

These are post commonly found in visceral sites such as gastrointestinal and uterus, in retroperitoneum and abdominal pelvic sites, rarely affecting skeletal system.

Case Report: A 45-year-old female was presented in the orthopedic outpatient department with a 2-year history of pain and swelling in her right hip with recent weight-bearing difficulties for 1 week after trivial slip and fall. X-rays revealed a lytic lesion in her right proximal femur along with neck of femur fracture. Biopsy revealed perivascular epithelioid tumor. Diagnosed with primary bone PEComa without metastasis in the right proximal femur, the patient underwent wide local excision and reconstruction using a mega-prosthesis.

Post-operative histopathological evaluation surprisingly identified a low-grade chondrosarcoma, but with clear surgical margins.

Conclusion: This case highlighted an unusual diagnostic discrepancy between pre-operative and post-operative evaluations. Nonetheless, the adopted strategy was effective, allowing the patient to resume daily activities. Surgeons should remain aware of the potential for primary bone tumors to masquerade as atypical tumors.

Keywords: Chondrosarcoma, perivascular epithelioid tumor, proximal femur, biopsy.

Introduction

Chondrosarcoma is the second most common primary malignant tumor of bone accounting for 20–27% of these [1]. Chondrosarcoma is malignant tumor; that produces a chondrosarcoma cartilaginous matrix [2]. Classification of chondrosarcoma is based on histological subtypes, including conventional (grades 1–3), clear cell, mesenchymal, and

dedifferentiated 2020 [3].

The World Health Organization classification divides chondrosarcoma into eight subtypes [4]. These are found most commonly, in the bones of the axial skeleton (pelvis, scapula, sternum, and ribs), followed by the proximal femur and proximal humerus [5]. The treatment for proximal femur tumor necessitates a two-part strategy incorporating tumor resection

Author's Photo Gallery



Dr. Ananya Mishra



Dr. Pulkit Gupta



Dr. Manohar



Dr. Hoysala Kantharaju



Dr. Dhyaan Bhardwaj



Dr. Vipul Jhajharia

Access this article online

Website:
www.jocr.co.in

DOI:
<https://doi.org/10.13107/jocr.2026.v16.i09.8076>

¹Department of Orthopaedics, AIIMS, New Delhi, India,
²Department of Orthopaedics, Sports Injury Centre, New Delhi, India,
³Department of Orthopaedics, SMS Medical College, Jaipur, India.

Address of Correspondence:

Dr. Ananya Mishra,
Department of Orthopaedics, AIIMS, New Delhi, India.
E-mail: mishraana.1308@gmail.com

Submitted: 14/06/2026; Review: 09/07/2026; Accepted: August 2026; Published: September 2026

DOI: <https://doi.org/10.13107/jocr.2026.v16.i09.8076>

© The Author(s). 2026 Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and non-commercial reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.



Figure 1: Radiograph of bilateral hip with pelvis of patient at presentation preoperatively showing a lytic lesion at right proximal femur with fracture neck of femur.

for oncological reasons and bone and soft-tissue reconstruction for mechanical purposes [6]. Perivascular epitheloid cell tumor (PEComa) is a rare mesenchymal neoplasm. These tumors all share a distinctive cell type, the perivascular epitheloid cell (which has no known normal tissue counter-part). These are most commonly found in visceral sites such as gastrointestinal and uterus, in retroperitoneum and abdominal pelvic sites [7], rarely affecting skeletal system.

Case Report

A 45-year-old female was presented in the orthopedic outpatient department with a 2 year history of pain and swelling in her right hip with recent weight-bearing difficulties for 1 week after trivial slip and fall. The patient was unable to bear full weight on affected limb since 1 week. She faced pain at hip joint in squatting and sitting cross legged since 2 year which were gradually increasing in intensity. There was history of weight loss. There was no association with fever, chills, or history of steroid intake. Radiograph (Fig. 1) revealed a lytic lesion in her right femur, involving the entire peri-trochanteric region, whole neck, parts of femur head, and proximal shaft which was compounded by a femur neck fracture.

Following initial management with traction and immobilization, further investigations, including magnetic resonance imaging (MRI) and positron emission tomography (PET) scan, were conducted. MRI (Fig. 2 and 3) revealed a T2 hyperintense, and T1 hypointense, mass confined to the bone sparing surrounding soft tissues and neurovascular structures. PET scan was negative for metastasis. A core-needle biopsy indicated a perivascular epitheloid tumor. Patient was diagnosed with primary bone PEComa without metastasis in the right proximal femur. Under regional anesthesia, the patient was kept in lateral decubitus position and southern approach was used to expose lesion. The patient underwent wide local excision and reconstruction using a mega-prosthesis. Post-operative radiograph (Fig. 4) showed complete excision of lytic area replaced with mega-prosthesis appropriately. The excised tissue was postoperatively sent for biopsy to confirm the diagnosis. The biopsy result surprisingly came out as low-grade chondrosarcoma showing presence of atypical chondrocytes with hyperchromatic nuclei and calcification (Fig. 5).

Discussion

The case report shows that sometime a common bone tumor can have atypical presentation.

Biopsy of lesion although being gold standard diagnostic tool cannot be 100% reliable due to inadequate tissue sample, wrong tissue sample or maybe due to inexperienced pathologist. PEComas are a family of mesenchymal tumors that rarely arise as a primary bone tumor [8]. Our patient had involvement of proximal femur which is also a common location for chondrosarcoma. The age of 45 years also favors for chondrosarcoma as it is common in adults of 30–70 years of age.




Figure 2: Coronal magnetic resonance imaging of bilateral hip and pelvis showing lytic lesion in proximal femur and fracture neck of femur right side.




Figure 3: Sagittal view of magnetic resonance imaging of bilateral hip with pelvis showing extent of lytic lesion in anteroposterior plane in the right proximal femur.

Clinically, most patients present with pain, swelling, and the presence of a mass of variable duration [9]. Radiologically, chondrosarcoma shows lytic lesions (50%) intralesional calcifications: ~70% (ring and arc or popcorn calcification), endosteal scalloping affecting more than two-thirds of the cortical thickness, moth-eaten appearance, or permeative appearance in higher grade tumors, for example, myxoid, dedifferentiated, and mesenchymal chondrosarcoma. Cortical remodeling and periosteal reaction are also seen. MRI shows low-to-intermediate signal intensity lesion on T1 and T2 shows high-signal intensity lesion as was consistent with our patient

[10]. Diagnosis is confirmed on biopsy. Macroscopically chondrosarcoma presents as translucent lobular, blue-gray, or white cut surface corresponding to the presence of hyaline cartilage. There may be areas containing myxoid or mucoid material and cystic changes. Microscopically, chondrosarcoma show abundant blue-gray cartilage matrix-production. Irregularly shaped lobules of cartilage varying in size and shape are present. Fibrous bands separate that these lobules or permeate bony trabeculae calcified areas suggesting the presence of a pre-existing enchondroma can often be found. The chondrocytes are atypical, with variable size and shape and contain enlarged hyperchromatic nuclei. Low-grade chondrosarcoma often closely resembles normal cartilage or the benign enchondroma, it is differentiated by presence of “chondrosarcoma permeation pattern,” where the tumor infiltrates through the marrow cavity instead of being confined by the native architecture as present in benign pathology [11]. Low grade or grade 1 chondrosarcoma with no aggressive imaging features can be treated with intralesional curettage and local adjuvants [12]. However, when aggressive biologic behavior is evident on imaging, wide resection following surgical principles of malignant bone tumors seems more appropriate. Our patient underwent wide resection and reconstruction with mega-prosthesis as the pre-operative biopsy diagnosis was PEComa which needs wide surgical excision [13]. However, despite the inconsistency in diagnosis, the patient had optimum final outcome.

Conclusion

This case highlighted an unusual diagnostic discrepancy between pre-operative and post-operative evaluations. Nonetheless, the adopted strategy was effective, allowing the patient to resume daily activities. Surgeon should always send



Figure 4: Post-operative radiograph of patient showing removal of lytic lesion and replacing proximal femur with mega prosthesis.




Figure 5: (a and b) Post-operative histopathological report showing grade 1 chondrosarcoma.

post-operative tumor sample for confirmation of diagnosis and if there is discrepancy in diagnosis do the needful for the well-being of the patient.

Clinical Message

Surgeons should remain aware of the potential for primary bone tumors to masquerade as atypical tumors.

Declaration of patient consent: The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient has given the consent for his/ her images and other clinical information to be reported in the journal. The patient understands that his/ her names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Conflict of interest: Nil **Source of support:** None

References

1. Murphey MD, Walker EA, Wilson AJ, Kransdorf MJ, Temple HT, Gannon FH. From the archives of the AFIP: Imaging of primary chondrosarcoma: Radiologic-pathologic correlation. *Radiographics* 2003;23:1245-78.
2. Marco RA, Gitelis S, Brebach GT, Healey JH. Cartilage tumors: Evaluation and treatment. *J Am Acad Orthop Surg* 2000;8:292-304.
3. Aigner T. Towards a new understanding and classification of chondrogenic neoplasias of the skeleton--biochemistry and cell biology of chondrosarcoma and its variants. *Virchows Arch* 2002;441:219-30.
4. Choi JH, Ro JY. The 2020 WHO classification of tumors of soft tissue: Selected changes and new entities. *Adv Anat Pathol* 2021;28:44-58.
5. Thorkildsen J, Taksdal I, Bjerkehaugen B, Haugland HK, Børge Johannesen T, Viset T, et al. Chondrosarcoma in Norway 1990-2013; An epidemiological and prognostic observational study of a complete national cohort. *Acta Oncol* 2019;58:273-82. Erratum in: *Acta Oncol* 2019;58:393.
6. Li B, Yu Y, Bao Y, Song J. Proximal femoral tumor resection followed by joint prosthesis replacement: A systematic review and meta-analysis. *BMC Musculoskelet Disord* 2023;24:779.
7. Hornick JL, Fletcher CD. PEComa: What do we know so far? *Histopathology* 2006;48:75-82.
8. Zhong J, Hu Y, Si L, Xing Y, Geng J, Jiao Q, et al. Primary perivascular epithelioid cell tumor (PEComa) in bone: A review of the literature and a case arising in the humerus with multiple metastases. *J Bone Oncol* 2020;26:100336.
9. Van Oosterom AT, Dirix LY. Chondrosarcoma and other rare bone sarcomas. *Curr Opin Oncol* 1990;2:495-9.
10. Varma DG, Ayala AG, Carrasco CH, Guo SQ, Kumar R, Edeiken J. Chondrosarcoma: MR imaging with pathologic correlation. *Radiographics* 1992;12:687-704.
11. Limaie F, Davis DD, Sticco KL. Chondrosarcoma. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2025. Available from: <https://www.ncbi.nlm.nih.gov/books/nbk538132> [Last accessed on 15 Mar 2025].
12. Campanacci DA, Scoccianti G, Franchi A, Roselli G, Beltrami G, Ippolito M, et al. Surgical treatment of central grade 1 chondrosarcoma of the appendicular skeleton. *J Orthop Traumatol* 2013;14:101-7.
13. Czarnecka AM, Skoczylas J, Bartnik E, Świtaj T, Rutkowski P. Management strategies for adults with locally advanced, unresectable or metastatic malignant perivascular epithelioid cell tumor (PEComa): Challenges and solutions. *Cancer Manag Res* 2023;15:615-23.

Conflict of Interest: Nil

Source of Support: Nil

Consent: The authors confirm that informed consent was obtained from the patient for publication of this article

How to Cite this Article

Mishra A, Gupta P, Manohar, Kantharaju H, Bhardwaj D, Jhajharia V. A Rare Case of Masquerading Chondrosarcoma with Pathological Fracture of Neck of Femur – A Case Report. *Journal of Orthopaedic Case Reports* 2026 September;16(09):315-318.

